

مراجع

- Celeux, G. and Diebolt, J. (1985), “The SEM algorithm: A probabilistic teacher algorithm derived from the EM algorithm for the mixture problem,” *Comput. Statist. Quart.*, 2, 73-82.
- Gelfand, A. E. (2000), “Gibbs Sampling,” *Journal of American Statistical Association*, 95, 1300-1304.
- Guo, S. W., and Thompson, E. A. (1992), “A Monte Carlo method for combined segregation and linkage analysis,” *American Journal of Human Genetics*, 51, 1111-1126.
- Jorgensen, B. (1984), “The delta algorithm and GLIM,” *International statistical Review*, 52, 283-300.
- Meilijson, I. (1989), “A fast improvement to the EM algorithm on its own terms,” *Journal of the Royal Statistical Society B*, 51, 127-138.
- Meng, X. L., and Rubin, D. B. (1991), “Using EM to obtain asymptotic variance-covariance matrices: The SEM algorithm,” *Journal of American Statistical Association*, 86, 899-909.
- Meng, X. L., and Rubin, D. B. (1993), “Maximum Likelihood Estimation via the ECM algorithm: a general framework,” *Biometrika*, 80, 267-278.
- Rai, S. N., and Matthews, D. E. (1993), “Improving the EM algorithm,” *Biometrics*, 49, 587-591.
- Ruud, P. A. (1991), “Extensions of estimation methods using the EM algorithm,” *J Econometrics*, 49, 305-341.
- Sinha, D., Tanner, M. A, and Hall, W. J. (1994), “Maximization of the marginal likelihood of grouped survival data,” *Biometrika*, 81, 53-60.
- Wei, G. C. G., and Tanner, M. A. (1990), “A Monte Carlo implementation of the EM algorithm and the poor man’s data augmentation algorithms,” *Journal of American Statistical Association*, 85, 699-704.